



# Results from the SDSS-II Supernova Survey

Rick Kessler  
University of Chicago

ANL Seminar Dec 9, 2008


# Contents

- ★ Dark Energy Intro
- ★ SDSS Supernova Survey (2005-07)
- ★ SN Ia Rate versus Redshift
- ★ Hubble Diagram Analysis & Results  
(1st-year SDSS data + external)


# **Primary Motivation for Supernova Surveys:**

**measure expansion  
history of the Universe:  
in particular, the role of  
dark energy**


# Understanding Expansion History is Tricky



# Understanding Expansion History is Tricky



# Understanding Expansion History is Tricky



# Fun Facts About Dark Energy

- $\rho_{\Lambda} = 10^{-29} \text{ g/cm}^3$  everywhere.
- Earth volume contains 0.01g of dark energy.
- Dark energy increases Earth's orbit by  $0.1 \mu\text{m}$ ; Pluto's orbit is increased by  $1 \mu\text{m}$ .
- Gravity and dark energy roughly cancel for Milky-Way and Andromeda galaxies (but galaxy-cluster gravity wins)
- $\Omega_{\Lambda} = 0.7$  today
- $\Omega_{\Lambda}/\Omega_M \sim 2.3$  today (compare  $\Omega_{\gamma}/\Omega_M < 10^{-4}$ ).
- $\Omega_{\Lambda} = \Omega_M$  at  $z=0.3$  (3-4 billion years ago, assumes  $w=-1$ ).
- Undetectable in terrestrial experiments (so far).
- Nobody knows what dark energy (or dark matter) is.

# Expansion Basics

$$H(z)^2 = H_0^2 \sum_i \Omega_i (1+z)^{3(1+w)}$$

[ where  $w = p/\rho$ , and  $R = 1/(1+z)$  ]

# Expansion Basics

$$H(z)^2 = H_0^2 \sum_i \Omega_i (1+z)^{3(1+w)}$$

[ where  $w = p/\rho$ , and  $R = 1/(1+z)$  ]

Source of expansion	w	Evolution with z	$\Omega$ at z=0
Matter (dark, baryon, relic $\nu$ )	$v^2/c^2 \sim 0$	$\Omega_M(1+z)^3$	0.3
Radiation (CMB)	1/3	$\Omega_\gamma(1+z)^4$	$\sim 10^{-5}$
Cosmological constant (?)	-1	$\Omega_\Lambda =$ constant	0.7
Curvature	-1/3	$\Omega_k(1+z)^2$	< few %

# Methods to Measure $H(z)$

$$H(z)^2 = \sum_i \Omega_i (1+z)^{3(1+w)}$$

<b>Method</b>	<b>Difficulties</b>
brightness vs. redshift	Large dispersion in brightness. Evolution ? Dust ?
count galaxy clusters vs redshift.	Need to know cluster-mass selection function.
galaxy clustering; power spectrum or clumpiness	galaxy vs. dark matter clustering
Weak lensing	Systematics of galaxy-shear measurements


# Methods to Measure $H(z)$

$$H(z)^2 = \sum_i \Omega_i (1+z)^{3(1+w)}$$

Method	Difficulties
brightness vs. redshift for <b>SN Ia</b>	<del>Large dispersion in brightness.</del> Evolution ? Dust ?


Natural dispersion ~ factor of 2 :  
reduced to 15% after 'width-luminosity' correction  
(Phillips 1993)

# Hubble Diagram Basics




Expansion history depends on  $\Omega_\Lambda$  and  $\Omega_M$

# Hubble Diagram Basics



Expansion history depends on  $\Omega_\Lambda$  and  $\Omega_M$




$$\text{mag} = -2.5 \log(\mathcal{L} / 4\pi d_L^2).$$

$$d_L = (1+z) \int dz / H(z, \Omega_M, \Omega_\Lambda, w)$$



for flat universe.

$$\text{Distance modulus: } \mu = 5 \log(d_L / 10 \text{ pc})$$


# Hubble Diagram Basics




Expansion history depends on  $\Omega_\Lambda$  and  $\Omega_M$



# w-sensitivity with Supernova



# w-Quest with Supernova



**$w = -0.9$  gives  
4% variation  
from  $w = -1$**

# Surveys


1990s

Development & discovery phase (Hi-z, SCP).

Lightcurve quality limited by telescope time.

compilation from Riess et. al., AJ 607 (2004):

Calan Tololo, HZT, SCP, CfA, Higher-Z, ACS.



# Surveys

1990s


Development & discovery phase (Hi-z, SCP).  
Lightcurve quality limited by telescope time.

2000s

Much more telescope time → rolling searches & more passbands.  
(SNLS, ESSENCE, SDSS)


compilation from Riess et. al., AJ 607(2004):

Calan Tololo, HZT, SCP, CfA, Higher-Z, ACS.



SNLS 1st year sample (Astier 2005)

plus ~ 40 low-z SNe from literature



# Surveys

1990s


Development & discovery phase (Hi-z, SCP).  
Lightcurve quality limited by telescope time.

2000s

Much more telescope time → rolling searches & more passbands.  
(SNLS, ESSENCE, SDSS)


compilation from Riess et. al., AJ 607(2004):

Calan Tololo, HZT, SCP, CfA, Higher-Z, ACS.



SNLS 1st year sample (Astier 2005)

plus ~ 40 low-z SNe from literature



# Surveys

1990s


Development & discovery phase (Hi-z, SCP).  
Lightcurve quality limited by telescope time.

2000s

Much more telescope time → rolling searches & more passbands.  
(SNLS, ESSENCE, SDSS)


compilation from Riess et. al., AJ 607 (2004):

Calan Tololo, HZT, SCP, CfA, Higher-Z, ACS.



SNLS 1st year sample (Astier 2005)

plus ~ 40 low-z SNe from literature



SDSS survey  
fills gap & adds  
low-z SNe



# SN papers becoming “Methodology” papers as surveys contribute smaller fraction of total SNe Ia

- **Astier06**: SNLS contributes ~ 70 of 110
- **Kowalski 2008**:  
contributes 8 of 307 SNe Ia
- **SDSS 2009**: contributes 103 of 288

# Meet the SDSS-II Supernova Team

## The Sloan Digital Sky Survey-II Supernova Survey: Technical Summary **AJ 135, 338 (2008)**

Joshua A. Frieman,<sup>1,2,3</sup> Bruce Bassett,<sup>4,5</sup> Andrew Becker,<sup>6</sup> Changsu Choi,<sup>7</sup> David Cinabro,<sup>8</sup> Fritz DeJongh,<sup>1</sup> Darren L. Depoy,<sup>9</sup> Ben Dilday,<sup>2,10</sup> Mamoru Doi,<sup>11</sup> Peter M. Garnavich,<sup>12</sup> Craig J. Hogan,<sup>6</sup> Jon Holtzman,<sup>13</sup> Myungshin Im,<sup>7</sup> Saurabh Jha,<sup>14</sup> Richard Kessler,<sup>2,15</sup> Kohki Konishi,<sup>16</sup> Hubert Lampeitl,<sup>17</sup> John Marriner,<sup>1</sup> Jennifer L. Marshall,<sup>9</sup> David McGinnis,<sup>1</sup> Gajus Miknaitis,<sup>1</sup> Robert C. Nichol,<sup>18</sup> Jose Luis Prieto,<sup>9</sup> Adam G. Riess,<sup>17,19</sup> Michael W. Richmond,<sup>20</sup> Roger Romani,<sup>14</sup> Masao Sako,<sup>21</sup> Donald P. Schneider,<sup>22</sup> Mathew Smith,<sup>18</sup> Naohiro Takahashi,<sup>11</sup> Kouichi Tokita,<sup>11</sup> Kurt van der Heyden,<sup>5</sup> Naoki Yasuda,<sup>16</sup> Chen Zheng,<sup>14</sup> Jennifer Adelman-McCarthy,<sup>1</sup> James Annis,<sup>1</sup> Roberto J. Assef,<sup>9</sup> John Barentine,<sup>23,24</sup> Ralf Bender,<sup>25,26</sup> Roger D. Blandford,<sup>14</sup> William N. Boroski,<sup>1</sup> Malcolm Bremer,<sup>27</sup> Howard Brewington,<sup>24</sup> Chris A. Collins,<sup>28</sup> Arlin Crotts,<sup>29</sup> Jack Dembicky,<sup>24</sup> Jason Eastman,<sup>9</sup> Alastair Edge,<sup>30</sup> Edmond Edmondson,<sup>18</sup> Edward Elson,<sup>5</sup> Michael E. Eyler,<sup>31</sup> Alexei V. Filippenko,<sup>32</sup> Ryan J. Foley,<sup>32</sup> Stephan Frank,<sup>9</sup> Ariel Goobar,<sup>33</sup> Tina Gueth,<sup>13</sup> James E. Gunn,<sup>34</sup> Michael Harvanek,<sup>24,35</sup> Ulrich Hopp,<sup>25,26</sup> Yutaka Ihara,<sup>11</sup> Željko Ivezić,<sup>6</sup> Steven Kahn,<sup>14</sup> Jared Kaplan,<sup>36</sup> Stephen Kent,<sup>1,3</sup> William Ketzeback,<sup>24</sup> Scott J. Kleinman,<sup>24,37</sup> Wolfram Kollatschny,<sup>38</sup> Richard G. Kron,<sup>3</sup> Jurek Krziesiński,<sup>24,39</sup> Dennis Lamenti,<sup>40</sup> Giorgos Leloudas,<sup>41</sup> Huan Lin,<sup>1</sup> Daniel C. Long,<sup>24</sup> John Lucey,<sup>30</sup> Robert H. Lupton,<sup>34</sup> Elena Malanushenko,<sup>24</sup> Viktor Malanushenko,<sup>24</sup> Russet J. McMillan,<sup>24</sup> Javier Mendez,<sup>42</sup> Christopher W. Morgan,<sup>9,31</sup> Tomoki Morokuma,<sup>11,43</sup> Atsuko Nitta,<sup>24,44</sup> Linda Ostman,<sup>33</sup> Kaike Pan,<sup>24</sup> Constance M. Rockosi,<sup>45</sup> A. Kathy Romer,<sup>46</sup> Pilar Ruiz-Lapuente,<sup>42</sup> Gabrelle Saurage,<sup>24</sup> Katie Schlesinger,<sup>9</sup> Stephanie A. Snedden,<sup>24</sup> Jesper Sollerman,<sup>41,47</sup> Chris Stoughton,<sup>1</sup> Maximilian Stritzinger,<sup>41</sup> Mark SubbaRao,<sup>3</sup> Douglas Tucker,<sup>1</sup> Petri Vaisanen,<sup>5</sup> Linda C. Watson,<sup>9</sup> Shannon Watters,<sup>24</sup> J. Craig Wheeler,<sup>23</sup> Brian Yanny,<sup>1</sup> and Donald York<sup>3,15</sup>

<sup>1</sup>Center for Particle Astrophysics, Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, IL 60510.

<sup>2</sup>Kavli Institute for Cosmological Physics, The University of Chicago, 5640 South Ellis Avenue Chicago, IL 60637.

<sup>3</sup>Department of Astronomy and Astrophysics, The University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637.

<sup>4</sup>Department of Mathematics and Applied Mathematics, University of Cape Town, Rondebosch 7701, South Africa.

<sup>5</sup>South African Astronomical Observatory, P.O. Box 9, Observatory 7935, South Africa.

<sup>6</sup>Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195.

<sup>7</sup>Department of Astronomy, Seoul National University, Seoul, South Korea.

<sup>8</sup>Department of Physics, Wayne State University, Detroit, MI 48202.

<sup>9</sup>Department of Astronomy, Ohio State University, 140 West 18th Avenue, Columbus, OH 43210-1173.

<sup>10</sup>Department of Physics, University of Chicago, Chicago, IL 60637.

<sup>11</sup>Institute of Astronomy, Graduate School of Science, University of Tokyo 2-21-1, Osawa, Mitaka, Tokyo 181-0015, Japan.

<sup>12</sup>University of Notre Dame, 225 Nieuwland Science, Notre Dame, IN 46556-5670.

<sup>13</sup>Department of Astronomy, MSC 4500, New Mexico State University, P.O. Box 30001, Las Cruces, NM 88003.

<sup>14</sup>Kavli Institute for Particle Astrophysics & Cosmology, Stanford University, Stanford, CA 94305-4060.

<sup>15</sup>Enrico Fermi Institute, University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637.

<sup>16</sup>Institute for Cosmic Ray Research, University of Tokyo, 5-1-5, Kashiwanoha, Kashiwa, Chiba, 277-8582, Japan.

<sup>17</sup>Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218.

<sup>18</sup>Institute of Cosmology and Gravitation, Mercantile House, Hampshire Terrace, University of Portsmouth, Portsmouth PO1 2EG, UK.

<sup>19</sup>Department of Physics and Astronomy, Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218.

<sup>20</sup>Physics Department, Rochester Institute of Technology, 85 Lomb Memorial Drive, Rochester, NY 14623-5603.

<sup>21</sup>Department of Physics and Astronomy, University of Pennsylvania, 203 South 33rd Street, Philadelphia, PA 19104.

<sup>22</sup>Department of Astronomy and Astrophysics, The Pennsylvania State University, 525 Davey Laboratory,

+ Steve, Joe, Hal

# SDSS-II Supernova Survey: Sep 1 - Nov 30, 2005-2007 (1 of 3 SDSS-II projects for 2005-2008)

## **GOAL:**

**Few hundred high-quality  
type Ia SNe lightcurves in  
redshift range 0.05-0.4**

## **SAMPLING:**

**~300 sq deg in ugriz  
(3 million galaxies every  
two nights)**

## **SPECTROSCOPIC FOLLOW-UP:**

**HET, ARC 3.5m, MDM,  
Subaru, WHT, Keck, NTT,  
KPNO, NOT, SALT,  
Magellan, TNG**




# SDSS Data Flow

One full night collects 800 fields (ugriz per field)  $\Leftrightarrow$  200 GB

Each 'search' field is compared to a 2-year old 'template' field ... things that go "boom" are extracted for human scanning.

Ten dual-CPU servers at APO process g,r,i data (2400 fields) in ~ 20 hrs.

one raw g-field ( $0.15^\circ$ )





(can you find a confirmed SN Ia ?)

# SDSS Data Flow







One full night collects 800 fields (ugriz per field)  $\Rightarrow$  200 GB

one raw g-field ( $0.15^\circ$ )



# SDSS Manual Scanning


Manual Scanning 10/20/2005 03:55 PM

	search	template	subtr	
<b>g</b>				g (srch, tmplt, subtr)
<b>r</b>				r (srch, tmplt, subtr)
<b>i</b>				i (srch, tmplt, subtr)

**The 'Good'**  
 **$z=0.05$  : also followed  
by SNF and CSP**

[http://sdssdp47.fnal.gov/sdssn\\_data/handscan/manual\\_scan.php](http://sdssdp47.fnal.gov/sdssn_data/handscan/manual_scan.php) Page 1 of 1


# SDSS Manual Scanning



**The 'Good'**

**$z=0.05$  : also followed  
by SNF and CSP**

[http://sdssdp47.fnal.gov/sdssn\\_data/handscan/manual\\_scan.php](http://sdssdp47.fnal.gov/sdssn_data/handscan/manual_scan.php)






**The 'Bad'**

[http://sdssdp47.fnal.gov/sdssn\\_data/handscan/manual\\_scan.php](http://sdssdp47.fnal.gov/sdssn_data/handscan/manual_scan.php)

Page 1 of 1

# SDSS Manual Scanning

Manual Scanning 10/20/2005 03:55 PM






**The 'Good'**

**$z=0.05$  : also followed  
by SNF and CSP**

[http://sdssdp47.fnal.gov/sdssn\\_data/handscan/manual\\_scan.php](http://sdssdp47.fnal.gov/sdssn_data/handscan/manual_scan.php)




Manual Scanning 10/20/2005 12:23 AM



**The 'Bad'**

[http://sdssdp47.fnal.gov/sdssn\\_data/handscan/manual\\_scan.php](http://sdssdp47.fnal.gov/sdssn_data/handscan/manual_scan.php)

Manual Scanning 10/20/2005



**The 'Ugly'**


[http://sdssdp47.fnal.gov/sdssn\\_data/handscan/manual\\_scan.php](http://sdssdp47.fnal.gov/sdssn_data/handscan/manual_scan.php)

Page

Manual Scan


$z = 0.09$

10/20/2005 04:47 PM




g (srch, tmplt,

subtr)



r (srch, tmplt,

subtr)



i (srch, tmplt,


subtr)

search    template    subtr

Manual Scan


$z = 0.29$

10/20/2005 04:49 PM




g (srch, tmplt,

subtr)



r (srch, tmplt,

subtr)




i (srch, tmplt,

subtr)

Manual Scan


$z = 0.20$

10/20/2005 04:48 PM




g (srch, tmplt,

subtr)



r (srch, tmplt,

subtr)



i (srch, tmplt,


subtr)

search    template    subtr

Manual Scan


$z = 0.36$

10/20/2005 04:50 PM




g (srch, tmplt,

subtr)



r (srch, tmplt,

subtr)




i (srch, tmplt,

subtr)


# Lightcurve Fits Update in Real Time

*2 epochs*


SN Ia Fit  
 $\chi^2 = 1.8$



SN Ibc Fit  
 $\chi^2 = 39$




SN II Fit  
 $\chi^2 = 6.6$




*30 epochs*


SN Ia Fit  
 $\chi^2 = 8.8$



SN Ibc Fit  
 $\chi^2 = 352$




SN II Fit  
 $\chi^2 = 57$




# Lightcurve Fits Update in Real Time

*2 epochs*


SN Ia Fit  
 $\chi^2 = 1.8$



SN Ibc Fit  
 $\chi^2 = 39$







SN II Fit  
 $\chi^2 = 6.6$



**> 90% of  
photometric Ia  
candidates were  
spectroscopically  
confirmed to be  
SN Ia**


# Follow-up Spectral id



# Survey Scan Stats

Sako et al., AJ 135, 348 (2008)

	<b>2005</b>
Objects scanned	<b>140,000</b>
SN candidates	<b>11,400</b>
Candidates with $\geq 1$ spectra	<b>180</b>
Confirmed SN Ia	<b>130</b>
Probable SN Ia	<b>16</b>
Confirmed SN other (Ib, Ic, II)	<b>18</b>



# Survey Scan Stats

Sako et al., AJ 135, 348 (2008)


	2005	2006	2007	Total
Objects scanned	140,000	14,400	15,200	170,000
SN candidates	11,400	3,700	3,967	19,000
Candidates with $\geq 1$ spectra	180	267	289	736
Confirmed SN Ia	130	197	171	498
Probable SN Ia	16	15	21	52
Confirmed SN other (Ib, Ic, II)	18	37	38	93

**Plus ~ 1000 photometric SN Ia:**


**we have 300 host-galaxy redshifts and still observing ...**

# SN Fakes


Fake SN Ia were inserted into the images in real time to measure software & scanning efficiencies.



# SDSS-SN Redshift & Cadence



# SDSS-SN Redshift & Cadence






Temporal edge effects: SNe peak too early or too late. May relax cuts later.

# SDSS SN Ia Lightcurves @


**$z = 0.09$**

**$z = 0.19$**

**$z = 0.36$**



# SDSS Rate for SN Ia with $z < 0.12$ : Fall-2005 sample → Dilday et al., arXiv:0801.3297 Motivation: understand nature of SN progenitors




Contributions:

16 spectroscopically confirmed Ia  
(26 before cuts)


1 photometric-id  
with host spec-Z

# Unbinned Likelihood Fit




# Rate vs. Redshift

P.Ripoche (AAS poster, Jan 2008)




# SDSS SN Ia Rate: in progress



**statistics  
vs.  
systematics**

# SDSS SN Ia Rate: in progress




**statistics  
vs.  
systematics**

# SDSS-II Hubble Diagram Analysis: Samples Include


- SDSS, Fall-2005 (103)
- Low redshift from literature (33)
- SNLS published (~ 70)
- ESSENCE published (~ 60)
- HST (34)

**Paper to be submitted  
soon to ApJ ...**



# Supernova Photometry from Fit

(Holtzman et. al., 2008, accepted by ApJ)



FIT-DATA:

all images (few dozen  $\times$  ugriz)

FIT-MODEL:

galaxy + stars + SN + sky

FIT PROPERTIES:

gal + stars: same in every image

SN: variable in every image

gal + stars + SN: PSF-smearred

**NO PIXEL RE-SAMPLING !**

⇒ no pixel correlations

⇒ proper stat. errors

# Extensive Photometry Tests Include:

- Recover zero flux pre-explosion
- Recover star mags
- Recover flux from fake SN

# Analysis Overview

- Use both **MLCS2k2** & **SALT2** methods without retraining ==> use essentially as-is
- Make necessary & obvious improvements to implementation, but not to underlying method.
- Identify problems & evaluate systematic uncertainties.

# Analysis Overview

- Six sample-combinations

a) SDSS-only

b) SDSS + ESSENCE + SNLS

c) Nearby + SDSS

d) Nearby + SDSS + ESSENCE + SNLS

e) Nearby + SDSS + ESSENCE + SNLS + HST

f) Nearby + ESSENCE + SNLS **(compare to WV07)**

**SDSS is  
high-z  
sample**




**no nearby  
sample; SDSS  
is lowz anchor**




# Lightcurve Fit: Brief Reminder

- Fit data to **parametric model** (or template) to get shape and color.
- Use shape and color to “standardize” intrinsic luminosity.



# Lightcurve Fit: Brief Reminder

- Fit data to **parametric model** (or template) to get shape and color.
- Use shape and color to “standardize” intrinsic luminosity.




## Jargon:

### "SN Training"

is the procedure for determining the relation between shape+color and intrinsic luminosity

# Lightcurve Fit: Brief Reminder


- Fit data to **parametric model** (or template) to get shape and color.
- Use shape and color to “standardize” intrinsic luminosity.



**Distance-modulus ( $\mu$ ) =**  
**Observed mag –**  
**Intrinsic mag**


# Lightcurve Fit: Brief Reminder

- Fit data to **parametric model** (or template) to get shape and color.
- Use shape and color to “standardize” intrinsic luminosity.





**Distance-modulus ( $\mu$ ) =**  
**Observed mag –**  
**Intrinsic mag –**  
**Dust-Extinction(??)**

# SN Ia Hubble Diagram





# Cosmology Fit for $w$ and $\Omega_M$ using SN+BAO+CMB



Preliminary


# Results



# Results

**ESSENCE: Wood-Vasey, AJ 666, 694 (2007)**


**SNLS: Astier, AJ 447, 31 (2006)**



# Questions to Ponder



- Q1: Why is SDSS-only systematic error so large ?
- Q2: Why does our MLCs-based  $w$ -result differ by  $\sim 0.3$  compared to WV07 (same method & same data) ?
- Q3: Why do MLCs and SALT2 results differ when high-redshift samples (ESSENCE + SNLS) are included ?

# Q1: Why is SDSS-only Systematic Error so Large ?




$$\sigma_w(\text{syst}) = .3$$
$$= 2 \times \sigma_w(\text{stat})$$

# Q1: Why is SDSS-only Systematic Error so Large ?




# Q1: Why is SDSS-only Systematic Error so Large ?



**Global SALT2  
“minimization”  
dampens U-bias:**

$$\sigma_w(\text{syst}) = .15$$

**Q2:** Why does our MLCS-based  $w$ -result differ by  $\sim 0.3$  compared to WV07 (same method & same data) ?



**Prior :**  
**survey**  
**efficiencies**  
**&**  
**host-galaxy**  
**dust**

# Comments on MLCS Fit Prior

- ❁ Prior 'P' = (underlying extinction distribution) x (observation efficiency):  $P=0$  when  $A_V < 0$
- ❁  $\chi^2 \rightarrow \chi^2 - 2\ln P$  (in light curve fit)
- ❁ Equivalent to using flat prior, and then applying bias-correction on each fitted distance modulus
- ❁ Either method (above) requires detailed simulation and knowledge of dust properties.

# Spectroscopic Inefficiency




Fig. 8.— Comparison of redshift distributions for data (dots) and simulations (histograms) for the four SN samples: nearby (Jha et al. 2007), SDSS-II (2005 sample), ESSENCE (Wood-Vasey et al. 2007), and SNLS (Astier et al. 2006). The nearby sample is shown on a loga-

- **Simulate all known effects using REAL observing conditions**
- **Compare data/sim redshift distributions**
- **Difference attributed to spectroscopic ineff.**

# Spectroscopic Inefficiency


(ignored in WV07)

- ★ Spectroscopic efficiency modeled as  $\exp(-m_V/\tau)$
- ★ Eff(spec) is included in fitting prior ...
- ★ SDSS-only w-shift is 0.13 ;  
much smaller w-shifts for sample-combinations
- ★ Assign w-syst error = 1/2 change from this effect

Dust Law:  $R_V = A_V/E(B-V)$   
and  $A(\lambda)$  from

Cardelli, Clayton, Mathis ApJ, 345, 245 (1989)

Blue light scatters more →  
extincted objects appear reddened.



$B + \Delta B$ ,  $V + \Delta V$   
→ earth

$$E(B - V) = \Delta B - \Delta V$$


# Dust Law: $R_V = A_V/E(B-V)$ and $A(\lambda)$ from

Cardelli, Clayton, Mathis ApJ, 345, 245 (1989)

- ★ Previous MLCS-based analyses assumed  $R_V = 3.1$  (global parameter)
- ★ Growing evidence points to  $R_V \sim 2$ :
  - SALT2 “ $\beta$ ” ( $R_V+1$ ) = 2 - 2.5
  - LOWZ studies (Nobili 08:  $R_V = 1.8$ )
  - individual SN with NIR (Krisciunas)
- ★ We have evaluated  $R_V$  with our own SDSS-II data

# Dust Law: $R_V = A_V/E(B-V)$


**To measure a global property of SN Ia,  
need sample with well-understood efficiency**



**Spec-confirmed SN Ia sample  
has large (spec) inefficiency  
that is not modeled by the sim.**


# Dust Law: $R_V = A_V/E(B-V)$

**To measure a global property of SN Ia,  
need sample with well-understood efficiency**




**Spec-confirmed SN Ia sample  
has large (spec) inefficiency  
that is not modeled by the sim.**

**Solution: include photometric  
SNe Ia with host-galaxy redshift !**



# Dust Law: $R_V = A_V/E(B-V)$

**Method: minimize  
data-MC chi2 for  
color vs. epoch**




# Dust Law: $R_V = A_V/E(B-V)$

**Method: minimize data-MC chi2 for color vs. epoch**

**SDSS Result:**

$$R_V = 2.23 \pm 0.14_{\text{stat}} \pm 0.45_{\text{syst}}$$

**Consistent with SALT2  $\beta$  and other SN-based studies.**



**Preliminary**

**Q2:** Why does our MLCS-based w-result differ by  $\sim 0.3$  compared to WV07 (same method & same data) ?

★  $R_V = 2.2$  (instead of 3.1 in WV07)

★ Account for spectroscopic inefficiency (ignored in WV07)

... and not discussed in this talk:

★ different  $A_V$  distribution than in WV07

★ Different treatment of Bessell filters for nearby ( $z < 0.1$ ) sample

★ Fit in flux instead of mag (avoids bias from removing  $\text{SNR} < 5$  measurements)

## Q3: SALT2-MLCS

Discrepancy:  $\delta w \sim 0.2$  ?



Oh no, these  
two light curve  
fitters give  
inconsistent  
values for  $w$  !

# Comparison of Lightcurve Fit Methods

<b>property</b>	<b>MLCS2k2 (Jha 2007)</b>	<b>SALT2/SNLS (Guy07)</b>
rest-frame model	<i>U,B,V,R,I</i> mag vs. t	spectral surface vs. t
color variations	host-galaxy dust	dust + intrinsic (no assump)
Fitting prior	Extinction $A_V > 0$	none

# Comparison of Lightcurve Fit Methods


property	MLCS2k2 (Jha 2007)	SALT2 (Guy07)
rest-frame model	$U, B, V, R, I$ mag vs. $t$	spectral surface vs. $t$
color variations	host-galaxy dust	dust + intrinsic (no assump)
Fitting prior	Extinction $A_V > 0$	none
K-correction	warp composite SN Ia spectrum from Hsiao	not needed
distance modulus	Fit-param for each SN Ia	from global fit

# Comparison of Lightcurve Fit Methods


property	MLCS2k2 (Jha 2007)	SALT2 (Guy07)
rest-frame model	$U, B, V, R, I$ mag vs. $t$	spectral surface vs. $t$
color variations	host-galaxy dust	dust + intrinsic (no assump)
Fitting prior	Extinction $A_V > 0$	none
K-correction	warp composite SN Ia spectrum from Hsiao	not needed
distance modulus	Fit-param for each SN Ia	from global fit
Training	$z < .1$ : SN lum & shape SDSS: $R_V, A_V$	all SNe used in training
Fitter availability	wrote our own fitter with improvements & options	black box provided by J.Guy of SNLS
Training availability	requires highly trained chef	Turn-key code, but crucial SNLS spectra are private



# MLCS-SALT2 Comparisons



# MLCS-SALT2 Comparisons




# Summary

- ★ SN-cosmology analysis with first-season SDSS-II sample (103 SNe Ia) is complete; paper near completion.
- ★  $w$ -measurements are systematics-limited.
- ★ Found lots of issues and problems to address using SDSS-SN for retraining & analysis with full 3-year sample.
- ★ Ideas welcome !!!
- ★ Warning/complaint: we really need turn-key training codes to study discrepancies & systematic uncertainties.

# Extra Slides

# Fit Residuals



# Fit Residuals

Table 13: Hubble diagram fit-quality parameters for each sub-sample.

fit-quality parameter	Result for sample:				
	Nearby	SDSS-II	ESSENCE	SNLS	HST
$\chi^2_{\mu}$ (independent fits)	31.2	53.6	49.6	59.3	33.4
$N_{dof}$	30	100	53	59	31
$RMS_{\mu}$	0.19	0.14	0.23	0.23	0.27




Preliminary

**smaller  $\chi^2$  is due to inefficiency  
from spectroscopic targeting.**

# Total Error Contours


(stretch stat-contour along BAO+CMB axis)



# Hubble Anomaly

a.k.a “Hubble Bubble”

Conley et. al. astro-ph/0705.0367




$\mu_{\text{fit}}$  from data;  $\mu_{\text{calc}}$  calculated from concordance model

# Hubble Anomaly

a.k.a “Hubble Bubble”

Conley et. al. astro-ph/0705.0367



# Conclusions

- ★ Paper in preparation (with 99 SDSS SNe Ia)
- ★ side-by-side comparisons of MLCS vs. SALT2
- ★ SDSS “*photometric SNe Ia +  $z_{\text{host}}$* ” are used to measure dust properties (RV) ... important step toward using photo-SN in Hubble diagram, and quantifying survey efficiency.
- ★ SDSS SN with  $z < .15$  may help understand low- $z$  Hubble anomaly.
- ★ Need publicly available training codes to optimize training and evaluate systematic errors.
- ★ all SDSS-based analysis (fitter & sim) is publicly available now ... data available with paper.
- ★ Three-season SDSS SN survey is done. Still acquiring host-galaxy redshifts to improve measurement of dust properties and for more SN Ia on the Hubble diagram.